
Specialty Code Set Training Gynecology Peggy Stilley, CPC, CPMA, CPC-I, CPB, COBGC Director, ICD-10 Training and Education Specialty Code Set Training Gynecology

No part of this presentation may be reproduced or transmitted in any form or by any means (graphically, electronically, or mechanically, including photocopying, recording, or taping) without the expressed written permission of AAPC.





Preventive Medicine Encounter for routine gyn exam Use additional code:

for screening of human papillomavirus, if applicable (Z11.51) for screening vaginal pap smear, if applicable (Z12.72)
To identify acquired absence of uterus, if applicable (Z90.71)

Encounter for gynecological exam (general) (routine) with abnormal findings
 Use additional code to identify the abnormal finding
 Encounter for gynecological exam (general) (routine) without abnormal findings



Specialty Code Set Training Gynecology

Example

 During a patient's annual gynecologic exam, the provider noticed a breast lump. The provider performs a cervical pap smear and orders a diagnostic mammogram.

-Z01.411 Encounter for gynecologic examination (general) (routine) with abnormal findings

-N63 Unspecified lump in breast




Specialty Code Set Training Gynecology

Abnormal Pap Smears

- R87.6-
- · Documentation should include the location (cervix, vagina)
- Cytology findings
 ASCUS
 - LGSIL
 - HGSIL
 - Inadequate smear
 - Thin preparation
 - Unsatisfactory smear
 - Satisfactory smear but lacking transformation zone
 - Other abnormal cytological findings on specimens



Other Cytology	
• R87.8 Human Papillomavirus HPV DNA tests	
Code selection is based on site and severity	
Cervical dysplasia N87.0 Mild (CIN1) N87.1 Moderate (CIN II)	
N87.9 Dysplasia of cervix, unspecified Severe (CIN III – reported from D06)	
Vaginal Dysplasia N89.0 Mild (VAIN1) N89.1 Moderate (VAIN II)	
N89.3 Dysplasia of vagina, unspecified Severe (VAIN III - reported from D07.2)	
AAPC Specialty Code Set Training Gynecology	
4	
Example	
The patient returns to the office to review the abnormal	
cervical pap smear results. The results show she has LGSIL.	
 -R87.612 Low grade squamous intraepithelial lesion on cytologic smear of cervix (LGSIL) 	-
AAPC Specialty Code Set Training Gynecology	
Fibroids	
Noncancerous growths (benign)Code selection based on	
Anatomic siteType	
 Intramural – located within the wall of the uterus Subserosal – located under the peritoneal surface of the 	
uterus Submucous – located in the muscle under the endometrium of the uterus	
ure dierus	
AAPC Specialty Code Set Training Gynecology	



D25.0 Submucous leiomyoma of uterus D25.1 Intramural leiomyoma of uterus D25.2 Subserosal leiomyoma of uterus D25.9 Leiomyoma of uterus D25.9 Leiomyoma of uterus, unspecified

Example


- Operative findings from an exploratory laparotomy and supracervical hysterectomy include a 12 week sized fibroid with the largest measuring approximately 8 cm in the left fundal region. She had normal appearing tubes and ovaries.
 - D25.9 Leiomyoma of uterus, unspecified



Vaginitis	
 Inflammation caused by infection or imbalance of vaginal bacteria Symptoms include: 	
ItchingDischarge	
– Pain	
 Light vaginal bleeding 	
AAPC Specialty Code Set Training Gynecology	
Vaginitis Code selection categorized by Acute or chronic Type Infectious agent Bacterial Candidiasis Trichomonas	
May require 2 codes – watch instructional notes	
AAPC Specialty Code Set Training Gynecology	
Example	
A patient presents with pelvic pain and abdominal	
discharge. The provider diagnosed the patient with bacterial vaginosis.	
- N76.0 Acute vaginosis	
- B96.89 Other specified bacterial agents as the cause of	
diseases classified elsewhere	
AAPC Specialty Code Set Training Gynecology	

Lichen	
Characterized by patchy, white thinning skin	
Occurs most often on the vulva	
Code selection categorized by:	
- Type	
- Location	
 If type is documented without site report L90.0 All types of lichen occurring on the vulva are coded as N90.4 	
7 in types of noneit occurring of the valva are coded as 1150.4	
AAPC Specialty Code Set Training Gynecology	
(Tail)AAPC Specialty Code Set Training Gynecology	
Example	
 During a genital exam, the provider notices a bed of 	
lichen sclerosis on the vulva	
- N90.4 Leukoplakia of vulva	
AAPC Specially Code Set Training Gynecology	
(T))AAPC Specialty Code Set Training Gynecology	
Menorrhagia	
Excessive or prolonged bleeding occurring at regular intervals	
Symptoms include	
Heavy bleedingPassing clots	
Anemia due to blood loss Code selection characterized by:	
- Primary	
Climacteric Menopausal, premenopausal	
PostclimactericPostmenopausal	-
Pubertal (menses retained)	
AAPC Specialty Code Set Training Gynecology	

	ICD-10 Codes	
N92.0	Excessive and frequent menstruation with regular cycle	
N92.1	Excessive and frequent menstruation with irregular cycle	
N92.2	Excessive menstruation at puberty	
N92.3	Ovulation bleeding	
N92.4	Excessive bleeding in the premenopausal period	
N92.5	Other specified irregular menstruation	
N92- Exclu	udesI note for N95.0 post-menopausal bleeding E30.1 precocious puberty(menstruation)	
N95.0	Post-menopausal bleeding	
(P) A A T		
(T)AAI	Specialty Code Set Training Gynecology	
	Example	
A . OI	O and former of the former of the DOO former of the former of the DOO former of the	
	o-Gyn performs a hysteroscopy with D&C for a patient osed with menorrhagia.	
a.a.g.r	Will more magain	
-N92.	Excessive and frequent menstruation with regular cycle	
₽ AAT		
(T)AAI	Specialty Code Set Training Gynecology	
	Overion Cyate	
	Ovarian Cysts	
	st is a sac filled with liquid or semi-fluid material in or	
	nd the ovary.	-
	otoms include:	
	egular menses elvic pain	
	nusea, vomiting	
	east tenderness	
	essure on rectum or bladder	
(Y)AAI		
(F)AAI	Specialty Code Set Training Gynecology	



Ovarian Cysts

- Pseudomucinois

- · Code selection is categorized by: - Multilocular
 - Adherent
 - Chocolate
 - Corpus Luteum

 - EndometrialDevelopmental
 - Dermoid
 - Follicular
- Serous
- Tuberculous

- Retention

- OviductCause/ contributing factor



Specialty Code Set Training Gynecology

ICD-10 Codes

	N80.1	Endometriosis of ovary (chocolate)	
	N83.0	Follicular cyst of ovary	
	N83.1	Corpus luteum cyst	
	N83.20	Unspecified ovarian cysts	
	N83.29	Other specified cyst (retention cyst, simple cyst)	
	neoplastic of	d N83.29 Excludes I developmental ovarian cyst (Q50.1), ovarian cyst (D27-), polycystic ovarian syndrome (E28.2), Stein- syndrome (E28.2)	
	N83 Excludes2 – hydrosalpinx N70.1		
6	AAPC	Specialty Code Set Training Gynecology	

Example • The findings for a diagnostic laparoscopy cystectomy includes a large amount of cle (hemoperitoneum) and right side of the al approximately 800 cc. Right ovarian cyst found to be bleeding at the time of surger were normal in appearance. Uterus and le normal in appearance. -N83.20 Ovarian cyst (hemorrhagic) -K66.1 Hemoperitoneum	ot within the pelvis bdomen, t filled with clot was ry. Tubes bilaterally	
(Y)AAPC Specialty C	Code Set Training Gynecology	
Menopausal Disorde • Category N95	ers	
Postmenopausal bleeding		
Menopausal Climacteric StatePostmenopausal atrophic vaginitis		
i ostriciopausai atropine vaginius		
AAPC Specialty C	Code Set Training Gynecology	

N95.1 Menopausal and female climacteric states Use an additional code for associated symptoms Excludes I asymptomatic menopausal syndrome (Z78.0), symptoms associated with artificial menopause (E89.41), symptoms associated with premature menopause (E28.310) N95.2 Postmenopausal atrophic vaginitis N95.8 Other specified menopausal and perimenopausal disorders

Examples	
 A patient presents complaining of headache and night sweats. The OBG contributes her symptoms to menopause. 	
- N95.1 Menopausal and female climacteric states	
R61 Generalized hyperhydrosisR51 Headaches	
AAPC Specialty Code Set Training Gynecology	
Example	
During a pelvic exam, the provider documents the patient has vaginal dryness and thin mucosa with diffuse erythema. The patient is diagnosed with	
atrophic vaginitis. The provider prescribes topical hormone replacement therapy.	
-N95.2 Post menopausal atrophic vaginitis	
Specialty Code Set Training Gynecology	
Specially Code Set Training Gynecology	
Pelvic Pain	
• R10.2 Pelvic and perineal pain	
Common symptoms associated with multiple conditions	
Abdominal pain has more options based on:	
- Type - Site	

Specialty Code Set Training Gynecology

(T)AAPC

The patient presents to her OBG complaining of pelvic pain lasting three days. She states the pain started right after her period. She is not experiencing any pain on unriation or during sexual activity. The provider performs a Pap smear and orders unrianglist and pelvic US -R10.2 Pelvic and perineal pain Prolapse Frequent problem for women having multiple births Symptoms include: - Discontine, pain - Protrusion of organs from introllus - Pressure - Urinary incontinence - Using incontinence - Conclusion based on: - Six of the prolapse - Complete or incomplete (or the stage of the prolapse) Stage of Prolapse - Userus drops to upper portion of vagina (incomplete) - Second degree prolapse - Userus drops to lover portion of the vagina (incomplete) - Third degree prolapse - Userus drops to lover portion of the vagina (incomplete) - Third degree prolapse - Userus drops to lover portion of the vagina (incomplete) - Third degree prolapse - Userus drops to lover portion of the vagina (incomplete) - Third degree prolapse - Userus drops to lover portion of the vagina (incomplete)	Example	
Prolapse Frequent problem for women having multiple births Symptoms include: Discontinual processing and providence performs a Pap smear and orders urinallysis and pelvic US -R10.2 Pelvic and perineal pain Prolapse Frequent problem for women having multiple births Symptoms include: Discontinual Prosure - Unimary incontinence Code selection based on: - Site of the prolapse - Complete or incomplete (or the stage of the prolapse) Prosure or women for the prolapse - Uterus drops to upper portion of vagina (incomplete) Second degree prolapse - Uterus drops to upper portion of the vagina (incomplete) Third degree prolapse - Uterus drops to upper portion of the vagina (incomplete) Third degree prolapse - Uterus drops to upper portion of the vagina (incomplete) Third degree prolapse - Uterus drops vith cervix located at vaginal opening (complete)		
-R10.2 Pelvic and perineal pain Prolapse Frequent problem for women having multiple births Symptoms include: - Disconfort, pain - Prontation of organs from introllus - Pressure - Urinary incontinence - Code selection based on: - Sixe of the prolapse - Complete or incomplete (or the stage of the prolapse) First degree prolapse - Uterus drops to upper portion of vagina (incomplete) - Second degree prolapse - Uterus drops to upper portion of the vagina (incomplete) - Third degree prolapse - Uterus drops to very province of the vagina (incomplete) - Third degree prolapse - Uterus drops to very portion of the vagina (incomplete) - Third degree prolapse - Uterus drops to very portion of the vagina (incomplete) - Third degree prolapse - Uterus drops to very portion of the vagina (incomplete)	period. She is not experiencing any pain on urination or during sexual activity. The provider performs a Pap smear	
Prolapse Frequent problem for women having multiple births Symptoms include: Disconfort, pain Protrusion of organs from introlius Pressure - Uniany incontinence Code selection based on: Site of the prolapse - Complete or incomplete (or the stage of the prolapse) Secondy Code Sea Treating Tymocotogy Secondy Code Sea Treating Tymocotogy Second Code Sea Treating Tymocotogy Uterus drops to upper portion of vagina (incomplete) - Uterus drops to lumper portion of the vagina (incomplete) Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)	and orders unnarysis and pervic US	
Prolapse Frequent problem for women having multiple births Symptoms include: Disconfort, pain Protrusion of organs from introllus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) First degree prolapse - Uterus drops to upper portion of vagina (incomplete) Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete)	-R10.2 Pelvic and perineal pain	
Prolapse Frequent problem for women having multiple births Symptoms include: Disconfort, pain Protrusion of organs from introllus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) First degree prolapse - Uterus drops to upper portion of vagina (incomplete) Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops to lower portion of the vagina (incomplete)		
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym	AAPC Specialty Code Set Training Gynecology	
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym	#	
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym		
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym		
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym		
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym		
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym		
Frequent problem for women having multiple births Symptoms include: Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Specially Code Bet Trailing Cymicology Special Cymicology Special Cym	Prolapse	
Discomfort, pain Protrusion of organs from introitus Pressure Urinary incontinence Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Stage of Prolapse First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete)		
- Protrusion of organs from introitus - Pressure - Urinary incontinence • Code selection based on : - Site of the prolapse - Complete or incomplete (or the stage of the prolapse) Stage of Prolapse • First degree prolapse - Uterus drops to upper portion of vagina (incomplete) • Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) • Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)		
- Urinary incontinence Code selection based on: - Site of the prolapse - Complete or incomplete (or the stage of the prolapse) Stage of Prolapse First degree prolapse - Uterus drops to upper portion of vagina (incomplete) Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)		
Code selection based on: Site of the prolapse Complete or incomplete (or the stage of the prolapse) Stage of Prolapse First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete)		
Stage of Prolapse First degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)		
Stage of Prolapse First degree prolapse - Uterus drops to upper portion of vagina (incomplete) Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)		
Stage of Prolapse • First degree prolapse - Uterus drops to upper portion of vagina (incomplete) • Second degree prolapse - Uterus drops to lower portion of the vagina (incomplete) • Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)		
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 	AAPC Specialty Code Set Training Gynecology	
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 		
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 		
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 		
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 		
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 		
 First degree prolapse Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 		
 Uterus drops to upper portion of vagina (incomplete) Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete) 	Stage of Prolapse	
Second degree prolapse Uterus drops to lower portion of the vagina (incomplete) Third degree prolapse Uterus drops with cervix located at vaginal opening (complete)		
- Uterus drops to lower portion of the vagina (incomplete) • Third degree prolapse - Uterus drops with cervix located at vaginal opening (complete)		
- Uterus drops with cervix located at vaginal opening (complete)	- Uterus drops to lower portion of the vagina (incomplete)	
(complete)		
AAPC Specialty Code Set Training Gynecology		
(F) AAPC Specialty Code Set Training Gynecology		
	AAPC Specialty Code Set Training Gynecology	

ICD-10 Codes

N81.0	Urethrocele	N81.5	Vaginal enterocele
N81.10	Cystocele, unspecified	N81.6	Rectocele
N81.11	Cystocele, midline	Us (R15)	se additional code for fecal incontinence
N81.12	Cystocele, unilateral	N81.81	Perineocele
N81.2	Incomplete uterovaginal prolapse	N81.83	Weakening of pubocervical tissue
N81.3	Complete uterovaginal prolapse	N81.84	Weakening of rectovaginal tissue
N81.4	Uterovaginal prolapse, unspecified	N81.85	Cervical stump prolapse
Excludes I – genital prolapse complicating pregnancy (O34.5), prolapse of vaginal wall after hysterectomy (N99.3)		N81.89	Other female genital prolapse

		70.0		
(141)	Α	A	PC	

Specialty Code Set Training Gynecology

Coding Tips

- Coding for rectocele has instructional note to code any associated fecal incontinence
- Not uncommon for female patients to have more than one type of prolapse
- For uterine prolapse and cystocele, rectocele or enterocele, select code only for the uterine prolapse. (Excludes note for N81.10-N81.12, N81.5, and N81.6)



Specialty Code Set Training Gynecolog

Example

- A patient diagnosed with a third degree prolapse and rectocele undergoes a total vaginal hysterectomy and posterior colporrhaphy repair.
 - N81.3 Complete uterovaginal prolapse



	Urinary Tract Infections	
•	Urinary tract infections are usually caused by bacteria	
•	Symptoms of UTI – Dysuria –	
	- Frequency	
	- Urgency - Pressure	
•	Instructional note states to code also the infectious agent	
•	(B95-B97) — Contains two combination codes	
(F	AAPC Specialty Code Set Training Gynecology	
4		
	Example	
	_	
•	The patient presents with lower abdominal pain and burning with urination. She has suffered from UTI in the past. The	
	provider performs a urinalysis and urine culture. The culture	
	returns positive for group B streptococcus.	
	- N39.0 Urinary tract infection, site not specified	
	 B95.1 Streptococcus, group B, as the cause of diseases classified elsewhere. 	
	-	
T	AAPC Specialty Code Set Training Gynecology	
	Urinary Incontinence	
	Urinary incontinence is the loss of bladder control.	
	Symptoms include:	
	Minimal leakage to uncontrolled wetting Occurring with cough or sneeze	
•	- Occurring with cough or sneeze Code selection based on:	
	- Type	
	- Cause	

(T)AAPC

	10	CD-10 Codes Incontinence
	N39.3	Stress incontinence *
	N39.41	Urge incontinence *
	N39.42	Incontinence without sensory awareness
	N39.43	Post-void dribbling
	N39.44	Nocturnal enuresis
	N39.45	Continuous dribbling
	N39.46	Mixed incontinence (urge and stress)
	N39.490	Overflow incontinence
	N39.498	Other specified (total or reflex)
		39.41 code also overactive bladder ote listed under N39.3 and N39.4
	1420	
)AAPC	Specialty Code Set Training Gynecology
	which was presulted in a will have her and a local of from her PCI	Example sents with predominant urge incontinence, receded by stress incontinence. This has also urinary tract infection about once a month. We obtain a MESA symptom score, voiding diary systourethroscopy. We will obtain her records P for further evaluation and review.
₹)AAPC	Specialty Code Set Training Gynecology
		Cystitis
•		ed in Category N30
•	- Frequency	ue.
	 Hematuria 	
	Cloudy or foul sDiscomfort	melling urine
	Low grade feve	
•	Code selection i	s based on:
	 Acute or chroni Location or cau 	
		bee hematuria, default without hematuria
		e to use additional code to identify infectious agent (B95-B97)
	AAPC	Specialty Code Set Training Gynecology

	ICD-10 Codes		
N30.0	N30.00 Acute cystitis, without hematuria N30.01 Acute cystitis, with hematuria		
N30.1	N30.10 Interstitial cystitis, without hematuria (chronic)		
N30.2	N30.11 Interstitial cystitis, with hematuria (chronic) N30.20 Other chronic cystitis, without hematuria		
N30.3	N30.21 Other chronic cystitis, with hematuria N30.30 Trigonitis, without hematuria		
N30.4	N30.31 Trigonitis, with hematuria N30.40 Irradiation cystitis, without hematuria		
N30.8	N30.41 Irradiation cystitis, with hematuria N30.80 Other cystitis, without hematuria (abscess of bladder)		
N30.9	N30.81 Other cystitis, with hematuria N30.90 Cystitis, unspecified without hematuria		
	N30.91 Cystitis, unspecified with hematuria		
Use addition	al code to identify infectious agent (B95-B97)		
AAPC	Specialty Code Set Training Gyne	cology	
	Example		
	presents with burning and frequency of urination. The		
	urinalysis and UA culture. The culture returns positiv		
	provider call in an antibiotic to treat the patient's acur	te	
cystitis.			
N30.00	Acute cystitis without hematuria		
B96.20	Unspecified Escherichia coli (E coli) as the cause of		
	s classified elsewhere		
AADO	THE RESERVE OF THE PARTY OF THE		
AAPC	Specialty Code Set Training Gyne	cology	
	Complications of Care Codes		
	·		
Intraop	erative and post-procedural complications are	Э	
	within the body system chapters with codes sp		
to the	organs and structures of that body system.	200110	
	ications codes are sequenced as the first-list	ed,	
followe	d by the code for the specific complication.		
AAPC	Specialty Code Set Training Gyne	cology	

N99.61	Intraoperative hemorrhage and hematoma of a genitourinary system organ or stricture	
N99.71	Accidental puncture and laceration of genitourinary organ system	
N99.72	Accidental puncture and laceration of genitourinary system, organ, structure during other procedure	
N99.81	Other intraoperative complication of genitourinary system	
N99.820	Post-procedural hemorrhage and hematoma of genitourinary system, organ, or structure following a genitourinary system procedure	
N99.821	Post-procedural hemorrhage and hematoma of genitourinary system, organ, or structure following an genitourinary system procedure	

