



ICD-9-CM Code	ICD-10-CM Code(s)	
<b>786.50</b>	Chest pain, unspecified	
ICD-10-CM Codes	<b>R07.9</b> Chest pain, unspecified **There are more specific code choice selections available in ICD-10-CM. These include: <b>R07.89</b> Other chest pain <b>R07.2</b> Precordial pain <b>R07.1</b> Chest pain on breathing <b>R07.81</b> Pleurodynia	
<b>414.00</b>	Coronary atherosclerosis of unspecified type of vessel, native or graft	
ICD-10-CM Codes	No map (No ICD-10 code exists for unspecified vessel; native or bypass graft must be indicated)	
<b>427.31</b>	Atrial fibrillation	
ICD-10-CM Codes	<b>I48.0</b> Paroxysmal atrial fibrillation <b>I48.1</b> Persistent atrial fibrillation <b>I48.2</b> Chronic atrial fibrillation <b>I48.91</b> Unspecified atrial fibrillation	
<b>427.89</b>	Cardiac dysrhythmias, other specified	
ICD-10-CM Codes	<b>I49.8</b> Other specified cardiac arrhythmias *Instructional note in ICD-10-CM indicates to code first cardiac arrhythmia complicating: abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) obstetric surgery and procedures ( <b>O75.4</b> )	
<b>414.01</b>	Coronary atherosclerosis of native coronary artery	
ICD-10-CM Codes	<b>I25.10</b> Atherosclerotic heart disease of native coronary artery without angina pectoris **Assign code from <b>I25.110-I25.119</b> when angina pectoris is present *Instructional note in ICD-10-CM indicates to use additional code, if applicable, to identify coronary atherosclerosis due to lipid rich plaque ( <b>I25.83</b> ) Use additional code to identify: chronic total occlusion of coronary artery ( <b>I25.82</b> ) exposure to environmental tobacco smoke ( <b>Z77.22</b> ) history of tobacco use ( <b>Z87.891</b> ) occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) tobacco dependence ( <b>F17-</b> ) tobacco use ( <b>Z72.0</b> )	
<b>794.31</b>	Abnormal electrocardiogram	<b>R94.31</b> Abnormal electrocardiogram
<b>424.1</b>	Aortic valve disorders	
ICD-10-CM Codes	<b>I35.9</b> Nonrheumatic aortic valve disorder, unspecified <b>I35.0</b> Nonrheumatic aortic (valve) stenosis <b>I35.1</b> Nonrheumatic aortic (valve) insufficiency <b>I35.2</b> Nonrheumatic aortic (valve) stenosis with insufficiency <b>I35.8</b> Other nonrheumatic aortic valve disorders	
<b>786.05</b>	Shortness of breath	<b>R06.02</b> Shortness of breath

ICD-9-CM Code	ICD-10-CM Code(s)	
<b>427.69</b>	Premature beats, NEC	
ICD-10-CM Codes	<b>I49.40</b> Unspecified premature depolarization <b>I49.1</b> Atrial premature depolarization <b>I49.2</b> Junctional premature depolarization <b>I49.3</b> Ventricular premature depolarization <b>I49.8</b> Other specified cardiac arrhythmias <b>I49.49</b> Other premature depolarization *Instructional note in ICD-10-CM indicates to code first cardiac arrhythmia complicating: abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) obstetric surgery and procedures ( <b>O75.4</b> )	
<b>427.1</b>	Paroxysmal ventricular tachycardia	<b>I47.2</b> Ventricular tachycardia
<b>425.4</b>	Primary cardiomyopathies, NEC	
ICD-10-CM Codes	<b>I42.2</b> Other hypertrophic cardiomyopathy <b>I42.5</b> Other restrictive cardiomyopathy <b>I42.8</b> Other cardiomyopathies <b>I42.9</b> Cardiomyopathy, unspecified (familial)(idiopathic) *Instructional note in ICD-10-CM indicates to code first cardiomyopathy complicating pregnancy and puerperium ( <b>O99.4</b> )	
<b>780.2</b>	Syncope and collapse	<b>R55</b> Syncope and collapse
<b>427.81</b>	Sinoatrial node dysfunction	
ICD-10-CM Codes	<b>I49.5</b> Sick sinus syndrome *Instructional note in ICD-10-CM indicates to code first cardiac arrhythmia complicating: abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) obstetric surgery and procedures ( <b>O75.4</b> )	
<b>786.09</b>	Dyspnea and respiratory abnormalities, other	<b>R06.89</b> Other abnormalities of breathing
<b>424.0</b>	Mitral valve disorders	
ICD-10-CM Codes	<b>I34.9</b> Nonrheumatic mitral valve disorder, unspecified <b>I34.0</b> Nonrheumatic mitral (valve) insufficiency <b>I34.1</b> Nonrheumatic mitral (valve) prolapse <b>I34.2</b> Nonrheumatic mitral (valve) stenosis <b>I34.8</b> Other nonrheumatic mitral valve disorders	
<b>794.30</b>	Abnormal cardiovascular function study, unspecified	
ICD-10-CM Codes	<b>R94.30</b> Abnormal result of cardiovascular function study, unspecified <b>R94.39</b> Abnormal result of other cardiovascular function study	
<b>427.9</b>	Cardiac dysrhythmia, unspecified	
ICD-10-CM Codes	<b>I49.9</b> Cardiac arrhythmia, unspecified ** Code first cardiac arrhythmia complicating: Abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) Obstetric surgery and procedures ( <b>O75.4</b> ) ** There are more specific code choice selections available in ICD-10-CM. These include: <b>I47.0</b> Re-entry ventricular arrhythmia <b>I47.1</b> Supraventricular tachycardia	



ICD-9-CM Code	ICD-10-CM Code(s)	ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont...)	<p><b>I47.2</b> Ventricular tachycardia  <b>I47.9</b> Paroxysmal tachycardia, unspecified          ** For <b>I47.0–I47.9</b> Code first cardiac arrhythmia complicating:          Abortion or ectopic or molar pregnancy (<b>O00-O07, O08.8</b>)          Obstetric surgery and procedures (<b>O75.4</b>)</p> <p><b>I48.0</b> Atrial fibrillation  <b>I48.1</b> Atrial flutter  <b>I49.01</b> Ventricular fibrillation  <b>I49.02</b> Ventricular flutter  <b>I49.1</b> Atrial premature depolarization  <b>I49.2</b> Junctional premature depolarization  <b>I49.3</b> Ventricular premature depolarization  <b>I49.40</b> Unspecified premature depolarization  <b>I49.49</b> Other premature depolarization  <b>I49.5</b> Sick sinus syndrome  <b>I49.8</b> Other specified cardiac arrhythmias          ** For <b>I49.0–I49.9</b> Code first cardiac arrhythmia complicating:          Abortion or ectopic or molar pregnancy (<b>O00-O07, O08.8</b>)          Obstetric surgery and procedures (<b>O75.4</b>)</p>	<b>401.9</b>	Hypertension (essential), unspecified
<b>785.1</b>	Palpitations	ICD-10-CM Codes	<p><b>I10</b> Essential (primary) hypertension          *Instructional note in ICD-10-CM indicates to use additional code to identify:          exposure to environmental tobacco smoke (<b>Z77.22</b>)          history of tobacco use (<b>Z87.891</b>)          occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)          tobacco dependence (<b>F17-</b>)          tobacco use (<b>Z72.0</b>)</p>
<b>ICD-10-CM Codes</b>	<b>R00.2</b> Palpitations <b>F45.8</b> Psychogenic palpitations	<b>426.0</b>	Atrioventricular block, complete
<b>429.3</b>	Cardiomegaly	<b>426.11</b>	Atrioventricular block, first degree
<b>428.22</b>	Systolic heart failure, chronic	<b>401.1</b>	Hypertension, benign
ICD-10-CM Codes	<p><b>I50.22</b> Chronic systolic (congestive) heart failure          *Instructional note in ICD-10-CM indicates to code first:          heart failure complicating abortion or ectopic or molar pregnancy (<b>O00-O07, O08.8</b>)          heart failure following surgery (<b>I97.13-</b>)          heart failure due to hypertension (<b>I11.0</b>)          heart failure due to hypertension with chronic kidney disease (<b>I13-</b>)          obstetric surgery and procedures (<b>O75.4</b>)          rheumatic heart failure (<b>I09.81</b>)</p>	ICD-10-CM Codes	<p><b>I10</b> Essential (primary) hypertension          *Instructional note in ICD-10-CM indicates to use additional code to identify:          exposure to environmental tobacco smoke (<b>Z77.22</b>)          history of tobacco use (<b>Z87.891</b>)          occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)          tobacco dependence (<b>F17-</b>)          tobacco use (<b>Z72.0</b>)</p>
<b>427.61</b>	Supraventricular premature beats	<b>272.4</b>	Hyperlipidemia, other and unspecified
ICD-10-CM Codes	<p><b>I49.1</b> Atrial premature depolarization          *Instructional note in ICD-10-CM indicates to code first cardiac arrhythmia complicating:          abortion or ectopic or molar pregnancy (<b>O00-O07, O08.8</b>)          obstetric surgery and procedures (<b>O75.4</b>)</p>	ICD-10-CM Codes	<p><b>E78.4</b> Other hyperlipidemia  <b>E78.5</b> Hyperlipidemia, unspecified          **There are more specific code choice selections available in ICD-10-CM. These include:  <b>E78.0</b> Pure hypercholesterolemia  <b>E78.1</b> Pure hyperglyceridemia  <b>E78.2</b> Mixed hyperlipidemia  <b>E78.3</b> Hyperchylomicronemia  <b>E78.6</b> Lipoprotein deficiency</p>
<b>427.0</b>	Paroxysmal supraventricular tachycardia	<b>V72.81</b>	Pre-operative cardiovascular exam
ICD-10-CM Codes	<p><b>I47.1</b> Supraventricular tachycardia          *Instructional note in ICD-10-CM indicates to code first tachycardia complicating:          abortion or ectopic or molar pregnancy (<b>O00-O07, O08.8</b>)          obstetric surgery and procedures (<b>O75.4</b>)</p>	<b>Z01.810</b>	Encounter for preprocedural cardiovascular examination
<b>786.59</b>	Chest pain, Other	<b>785.2</b>	Cardiac murmurs, undiagnosed
ICD-10-CM Codes	<b>R07.89</b> Other chest pain NOS <b>R07.82</b> Intercostal pain	<b>428.0</b>	Congestive heart failure, unspecified
		ICD-10-CM Codes	<p><b>I50.9</b> Heart failure, unspecified  <b>I50.1</b> Left ventricular failure  <b>I50.20</b> Unspecified systolic (congestive) heart failure  <b>I50.21</b> Acute systolic (congestive) heart failure  <b>I50.22</b> Chronic systolic (congestive) heart failure  <b>I50.23</b> Acute on chronic systolic (congestive) heart failure  <b>I50.30</b> Unspecified diastolic (congestive) heart failure  <b>I50.31</b> Acute diastolic (congestive) heart failure  <b>I50.32</b> Chronic diastolic (congestive) heart failure  <b>I50.33</b> Acute on chronic diastolic (congestive) heart failure  <b>I50.40</b> Unspecified combined systolic (congestive) and diastolic (congestive) heart failure  <b>I50.41</b> Acute combined systolic (congestive) and diastolic (congestive) heart failure</p>



ICD-9-CM Code	ICD-10-CM Code(s)	ICD-9-CM Code	ICD-10-CM Code(s)
<b>428.0</b> cont..	Congestive heart failure, unspecified	<b>428.42</b>	Combined systolic and diastolic heart failure, chronic
ICD-10-CM Codes cont...	<b>I50.42</b> Chronic combined systolic (congestive) and diastolic (congestive) heart failure <b>I50.43</b> Acute on chronic combined systolic (congestive) and diastolic (congestive) heart failure  *Instructional note in ICD-10-CM indicates to code first: heart failure complicating abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) heart failure following surgery ( <b>I97.13-</b> ) heart failure due to hypertension ( <b>I11.0</b> ) heart failure due to hypertension with chronic kidney disease ( <b>I13-</b> ) obstetric surgery and procedures ( <b>O75.4</b> ) rheumatic heart failure ( <b>I09.81</b> )	ICD-10-CM Codes	<b>I50.42</b> Chronic combined systolic (congestive) and diastolic (congestive) heart failure  *Instructional note in ICD-10-CM indicates to code first: heart failure complicating abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) heart failure following surgery ( <b>I97.13-</b> ) heart failure due to hypertension ( <b>I11.0</b> ) heart failure due to hypertension with chronic kidney disease ( <b>I13-</b> ) obstetric surgery and procedures ( <b>O75.4</b> ) rheumatic heart failure ( <b>I09.81</b> )
<b>414.8</b>	Other specified forms of chronic ischemic heart disease	<b>411.1</b>	Intermediate coronary syndrome
ICD-10-CM Codes	<b>I25.89</b> Other forms of chronic ischemic heart disease <b>I25.9</b> Ischemic heart disease; unspecified	ICD-10-CM Codes	<b>I20.0</b> Unstable angina  *Instructional note in ICD-10-CM indicates to use additional code to identify: exposure to environmental tobacco smoke ( <b>Z77.22</b> ) history of tobacco use ( <b>Z87.891</b> ) occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) tobacco dependence ( <b>F17-</b> ) tobacco use ( <b>Z72.0</b> )
<b>427.32</b>	Atrial flutter	<b>412</b>	Myocardial infarct, old
ICD-10-CM Codes	<b>I48.3</b> Typical atrial flutter <b>I48.4</b> Atypical atrial flutter <b>I48.92</b> Unspecified atrial flutter	ICD-10-CM Codes	<b>I25.2</b> Old myocardial infarction  *Instructional note in ICD-10-CM indicates to use additional code to identify: exposure to environmental tobacco smoke ( <b>Z77.22</b> ) history of tobacco use ( <b>Z87.891</b> ) occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) tobacco dependence ( <b>F17-</b> ) tobacco use ( <b>Z72.0</b> )
<b>426.4</b>	Right bundle branch block	<b>426.3</b>	Left bundle branch block, other and unspecified
ICD-10-CM Codes	<b>I45.10</b> Unspecified right bundle-branch block <b>I45.19</b> Other right bundle-branch block <b>I45.2</b> Bifascicular block	ICD-10-CM Codes	<b>I44.7</b> Left bundle-branch block, unspecified <b>I45.2</b> Bifascicular block <b>I44.60</b> Unspecified bifascicular block <b>I44.4</b> Left anterior bifascicular block <b>I44.5</b> Left posterior bifascicular block
<b>780.79</b>	Other malaise and fatigue	<b>426.50</b>	Bundle branch block, unspecified
ICD-10-CM Codes	<b>R53.81</b> Other malaise <b>R53.1</b> Weakness <b>R53.83</b> Other fatigue <b>G93.3</b> Postviral fatigue syndrome	ICD-10-CM Codes	<b>I45.4</b> Nonspecific intraventricular block <b>I44.30</b> Unspecified atrioventricular block <b>I44.39</b> Other atrioventricular block
<b>413.9</b>	Angina pectoris, other and unspecified	<b>425.11</b>	Hypertrophic obstructive cardiomyopathy
ICD-10-CM Codes	<b>I20.8</b> Other forms of angina pectoris <b>I20.9</b> Angina pectoris, unspecified <b>I20.0</b> Unstable angina <b>I20.1</b> Angina pectoris with documented spasm  *Instructional note in ICD-10-CM indicates to use additional code to identify: exposure to environmental tobacco smoke ( <b>Z77.22</b> ) history of tobacco use ( <b>Z87.891</b> ) occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) tobacco dependence ( <b>F17-</b> ) tobacco use ( <b>Z72.0</b> )	ICD-10-CM Codes	<b>I42.1</b> Obstructive hypertrophic cardiomyopathy  *Instructional note in ICD-10-CM indicates to code first cardiomyopathy complicating pregnancy and puerperium ( <b>O99.4</b> )
<b>780.4</b>	Dizziness and giddiness	<b>R42</b>	Dizziness and giddiness



ICD-9-CM Code	ICD-10-CM Code(s)	ICD-9-CM Code	ICD-10-CM Code(s)
<b>424.2</b>	Tricuspid valve disorders, specified as nonrheumatic	<b>416.8</b>	Pulmonary heart diseases, other, chronic
ICD-10-CM Codes	<b>I36.0</b> Nonrheumatic tricuspid (valve) stenosis <b>I36.1</b> Nonrheumatic tricuspid (valve) insufficiency <b>I36.2</b> Nonrheumatic tricuspid (valve) stenosis with insufficiency <b>I36.8</b> Other nonrheumatic tricuspid valve disorders <b>I36.9</b> Nonrheumatic tricuspid valve disorder, unspecified	ICD-10-CM Codes	<b>I27.2</b> Other secondary pulmonary hypertension <b>I27.89</b> Other specified pulmonary heart disease <b>I27.9</b> Pulmonary heart disease, unspecified <small>*Instructional note under <b>I27.2</b> states to code also associated underlying condition</small>
<b>410.40</b>	Myocardial infarction, acute, of other inferior wall, episode of care unspecified	<b>426.13</b>	Atrioventricular block, second degree, other
ICD-10-CM Codes	<b>I21.19</b> ST elevation (STEMI) myocardial infarction involving other coronary artery of inferior wall <b>I22.1</b> Subsequent ST elevation (STEMI) myocardial infarction of inferior wall <small>*Instructional note in ICD-10-CM indicates to use additional code to identify:</small> exposure to environmental tobacco smoke ( <b>Z77.22</b> ) history of tobacco use ( <b>Z87.891</b> ) occupational exposure to environmental tobacco smoke ( <b>Z57.31</b> ) status post administration of tPA (rtPA) in a different facility within the last 24 hours prior to admission to current facility ( <b>Z92.82</b> ) tobacco dependence ( <b>F17-</b> ) tobacco use ( <b>Z72.0</b> )	<b>782.3</b>	Edema
<b>785.0</b>	Tachycardia, unspecified	ICD-10-CM Codes	<b>R60.0</b> Localized edema <b>R60.1</b> Generalized edema <b>R60.9</b> Edema, unspecified
	<b>R00.0</b> Tachycardia, unspecified	<b>427.41</b>	Ventricular fibrillation
		ICD-10-CM Codes	<b>I49.01</b> Ventricular fibrillation <small>*Instructional note in ICD-10-CM indicates to code first cardiac arrhythmia complicating:</small> abortion or ectopic or molar pregnancy ( <b>O00-O07, O08.8</b> ) obstetric surgery and procedures ( <b>O75.4</b> )
		<b>416.0</b>	Primary pulmonary hypertension
			<b>I27.0</b> Primary pulmonary hypertension

*Fast Forward*  
to ICD-10



ICD-10